

Acute Effects of Heat-Not-Burn, Electronic Vaping, and Traditional Tobacco Combustion Cigarettes: The Sapienza University of Rome-Vascular Assessment of Proatherosclerotic Effects of Smoking (SUR-VAPES) 2 Randomized Trial

Giuseppe Biondi-Zoccai, MD, MStat;* Sebastiano Sciarretta, MD, PhD;* Christopher Bullen, MD; Cristina Nocella, PhD; Francesco Violi, MD; Lorenzo Loffredo, MD; Pasquale Pignatelli, MD; Ludovica Perri, MD; Mariangela Peruzzi, MD, PhD; Antonino G.M. Marullo, MD, PhD; Elena De Falco, PhD; Isotta Chimenti, PhD; Vittoria Cammisotto, MSc; Valentina Valenti, MD; Flaminia Coluzzi, MD; Elena Cavarretta, MD; Albino Carrizzo, PhD; Francesco Prati, MD; Roberto Carnevale, PhD;† Giacomo Frati, MD, MSc†

Background—Little clinical research on new-generation heat-not-burn cigarettes (HNBC) in comparison with electronic vaping cigarettes (EVC) and traditional tobacco combustion cigarettes (TC) has been reported. We aimed to appraise the acute effects of single use of HNBC, EVC, and TC in healthy smokers.

Methods and Results—This was an independent, cross-over, randomized trial in 20 TC smokers, with allocation to different cycles of HNBC, EVC, and TC. All participants used all types of products, with an intercycle washout of 1 week. End points were oxidative stress, antioxidant reserve, platelet activation, flow-mediated dilation, blood pressure, and satisfaction scores. Single use of any product led to an adverse impact on oxidative stress, antioxidant reserve, platelet function, flow-mediated dilation, and blood pressure. HNBC had less impact than EVC and TC on soluble Nox2-derived peptide (respectively, $P=0.004$ and 0.001), 8-isoprostaglandin F $_{2\alpha}$ -III ($P=0.004$ and <0.001), and vitamin E ($P=0.018$ and 0.044). HNBC and EVC were equally less impactful than TCs on flow-mediated dilation ($P=0.872$ for HNBC versus EVC), H $_2$ O $_2$ ($P=0.522$), H $_2$ O $_2$ breakdown activity ($P=0.091$), soluble CD40 ligand ($P=0.849$), and soluble P-selectin ($P=0.821$). The effect of HNBC and, to a lesser extent EVC, on blood pressure was less evident than that of TC, whereas HNBC appeared more satisfying than EVC (all $P<0.05$).

Conclusions—Acute effects of HNBC, EVC, and TC are different on several oxidative stress, antioxidant reserve, platelet function, cardiovascular, and satisfaction dimensions, with TCs showing the most detrimental changes in clinically relevant features.

Clinical Trial Registration—URL: <http://www.clinicaltrials.gov>. Unique identifier: NCT03301129. (*J Am Heart Assoc.* 2019;8:e010455. DOI: 10.1161/JAHA.118.010455.)

Key Words: flow-induced dilation • oxidative stress • platelet • platelet aggregation • smoking

It is estimated that cigarette smoking explains almost 90% of lung cancer risk in men and 70% to 80% in women and is responsible for \approx 140 000 premature deaths annually from

cardiovascular disease.^{1,2} Among the different harmful effects of smoking, oxidative stress plays an important role. Indeed, smoking may impair oxidative balance by inducing the

From the Department of Medico-Surgical Sciences and Biotechnologies, Sapienza University of Rome, Latina, Italy (G.B.-Z., S.S., M.P., A.G.M.M., E.D.F., I.C., V.V., F.C., E.C., R.C.); *Mediterranea - Cardiocentro*, Napoli, Italy (G.B.-Z., M.P., E.D.F., I.C., E.C., R.C.); IRCCS NEUROMED, Pozzilli, Italy (S.S., C.N., A.C., G.F.); National Institute for Health Innovation, School of Population Health, Faculty of Medical and Health Sciences, The University of Auckland, New Zealand (C.B.); Department of Internal Medicine and Medical Specialties, Sapienza University of Rome, Italy (F.V., L.L., P.P., L.P., V.C.); Division of Cardiology, San Giovanni Addolorata Hospital, Rome, Italy (F.P.); Centro Per La Lotta Contro L'Infarto, Rome, Italy (F.P.).

Accompanying Data S1, Table S1, and Figures S1 through S14 are available at <https://www.ahajournals.org/doi/suppl/10.1161/JAHA.118.010455>

*Dr Biondi-Zoccai and Dr Sciarretta equally contributed to this work.

†Dr Carnevale and Dr Frati are co-senior authors.

Correspondence to: Giuseppe Biondi-Zoccai, MD, MStat, Department of Medico-Surgical Sciences and Biotechnologies, Sapienza University of Rome, Corso della Repubblica 74, 04100 Latina, Italy. E-mail: giuseppe.biondizoccai@uniroma1.it

Received July 27, 2018; accepted December 18, 2018.

© 2019 The Authors. Published on behalf of the American Heart Association, Inc., by Wiley. This is an open access article under the terms of the Creative Commons Attribution-NonCommercial License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

Clinical Perspective

What Is New?

- Limited comparative research on new-generation heat-not-burn cigarettes, electronic vaping cigarettes, and traditional tobacco combustion cigarettes (TCs) has been reported.
- We performed a cross-over, randomized trial in 20 TC smokers, with allocation to different cycles of heat-not-burn cigarettes, electronic vaping cigarettes, and TCs, without any funding from tobacco or electronic vaping cigarette companies.

What Are the Clinical Implications?

- We found that acute effects of heat-not-burn cigarettes, electronic vaping cigarettes, and TCs are different on several oxidative stress, antioxidant reserve, platelet function, cardiovascular, and satisfaction dimensions, with TCs showing the most detrimental changes in clinically relevant features, thus suggesting that these modified risk products may indeed prove as a useful tool to quit TCs.

production of reactive oxygen species and by weakening the antioxidant defense systems. Moreover, smoking promotes atherothrombosis by inducing platelet activation, and we previously demonstrated that chronic smokers displayed different levels of biomarkers of oxidative stress and platelet activation compared with nonsmokers, and also responded differently to different types of cigarettes.^{3,4}

Electronic vaping cigarettes (EVCs), modern technological surrogates of traditional tobacco combustion cigarettes (TCs), were introduced in 2004.^{5,6} They are battery-operated aerosolizing devices using liquids that entail a mixture of glycerol and propylene glycol, flavors, and, most commonly, nicotine ranging from 1.6 to 19 mg/cartridge.⁷ EVCs are considered a relatively less harmful alternative to TCs given that no combustion occurs, and indeed several patent applications describe EVCs as electronic atomization cigarettes that contain only nicotine without tobacco tar.⁸ However, their long-term safety remains largely uncertain, especially in patients with established clinical disease.⁹

A new-generation heat-not-burn cigarette (HNBC) has been recently introduced.¹⁰ This device heats a disposable tobacco stick with a thin metallic blade. The stick is maintained at a controlled heating temperature up to 350°C, without combustion, fire, ash, or smoke. In addition, in contrast to EVCs, HNBCs do not vaporize liquid-containing flavorings, propylene glycol, and vegetable glycerol. However, this new device needs to be fully examined in clinical studies, particularly in the cardiovascular system.¹¹

Recently, we and others highlighted the acute adverse effects *in vivo* and *in vitro* of EVCs on vascular function,

oxidative stress, and platelet activation, which are recognized as pathophysiological factors for atherosclerosis development and progression and, ultimately, vascular disease.^{12–14} Specifically, we found that EVCs had less impact than TCs on oxidative stress, flow-mediated dilation (FMD), and platelet aggregation. Moreover, Nabavizadeh et al demonstrated that, in rats, acute exposures to HNBC aerosol impairs endothelial function, assessed by FMD.¹⁵ However, no independent comparative data on the acute impact of HNBCs on vascular function, oxidative stress, and platelet activation are available in chronic TC users.

We thus aimed at comparing HNBCs, EVCs, and TCs, focusing on acute effects on oxidative stress, FMD, platelet function, blood pressure, and satisfaction.

Methods

The data, analytical methods, and study materials of this trial will be made available to other researchers for purposes of reproducing the results or replicating the procedure, upon written request.

Design

This was an independent, randomized, cross-over study conducted in 2017 at Sapienza University of Rome, Latina, Italy. All participants provided written informed consent. The study was approved by the ethics committee of Sapienza University of Rome (06-27-2014, protocol number 813/14) and conducted in accord with the Declaration of Helsinki. No funding, directly or indirectly, was received from any HNBC, EVC, or TC manufacturer or supplier, or affiliated organizations.

Participants

We recruited apparently healthy subjects, on the basis of the following: (1) no acute or chronic organic, metabolic, and inflammatory disease; (2) no fever or infection within 3 months; (3) no cardiovascular symptoms; (4) no allergies; and (5) normal blood pressure levels and heart rhythm at screening. Women were not pregnant and not menstruating when the experiment was performed. In the month preceding the study and subsequently, none of the participants took vitamin E, other antioxidant supplements, or other drugs known to affect or modulate oxidative stress, FMD, platelet function, or blood pressure. A 1-week washout from any tobacco product was recommended at study entry and before each experimental cycle, and smoking history (time when smoking had begun) and intensity (ie, number of daily cigarettes) was self-reported.

Procedures

Participants were previously tutored on how to use the different devices correctly by means of standardized videos. They were then randomly allocated to 6 different cycles of using a single HNBC, EVC, and TC. Participants either smoked a single TC from a leading brand (Marlboro Gold; Philip Morris International, Neuchatel, Switzerland) with a mean nicotine content of 0.60 mg per cigarette according to the package label, vaped 9 puffs from a leading tobacco-flavored EVC (Blu Pro, Fontem, Netherlands), charged with a nicotine cartridge with a mean nicotine content of 16 mg, equivalent to 250 puffs according to the package label, thus yielding 0.58 mg of nicotine content in 9 puffs, or used a leading HNBC (THS2.2 IQOS with Heets; Amber Label, Philip Morris International) having a mean nicotine content of 0.50 mg per stick according to the manufacturer. Two investigators (who were past smokers or current users of all the devices) directly supervised all sessions to prevent incorrect product use or hyperventilation.

The randomization list was computer generated, and subjects were unaware of the assignment until each smoking session commenced. Participants eventually used all 3 products, with an intercycle washout period of 1 week. Blood samples were drawn just before and immediately after using each allocated product and analyzed for markers of oxidative stress, antioxidant reserve, platelet function, and cotinine levels. Endothelial dysfunction (as assessed by FMD) and cardiovascular parameters were analyzed, as previously reported.¹²

A 7-question product satisfaction questionnaire was administered after each smoking session, posing the following questions: “was the cigarette satisfying?”; “was the cigarette enjoyable?”; “did you experience a bad taste?”; “did you suffer from unexpected cough?”; “did smoking make you feel sick?”; “did smoking give you energy?”; and “soon after smoking did your desire for another cigarette decrease?”. Each feature was scored using a visual analogue scale, spanning from 0 (no effect) to 100 (maximum effect) according to Shiffman and Terhorst.¹⁶

All measurements were conducted by personnel blinded to the actual product used by the participant during each experimental session.

Laboratory Analysis

All materials for laboratory tests were from Sigma-Aldrich (St. Louis, MO), unless otherwise specified. Blood obtained from subjects was collected in blood collection tubes with or without anticoagulant and centrifuged at 300g for 10 minutes to obtain supernatant. Serum and plasma samples were instantly stored at -80°C until use with the antioxidant butylated hydroxytoluene at a final concentration of 20 mmol/L.

Ultrasound assessment of basal brachial diameter and endothelial-dependent FMD of the brachial artery were investigated according to current guidelines.^{16,17} Briefly, the study was performed in a temperature-controlled room (22.8°C) with subjects in a resting, supine state. All evaluations were performed between 8 and 10 AM; brachial artery diameter was imaged using a 7.5-MHz linear array transducer ultrasound system (Vivid S6; GE Healthcare, Little Chalfont, UK) equipped with electronic callipers, vascular software for 2-dimensional imaging, color and spectral Doppler, and internal ECG. The brachial artery was imaged at a location 3 to 7 cm above the antecubital crease. In order to create a flow stimulus in the brachial artery, a sphygmomanometric cuff was placed on the forearm. The cuff was inflated at least 50 mm Hg above systolic pressure to occlude artery inflow for 5 minutes. All vasodilatation measurements were made at the end of diastole. FMD was expressed as change in poststimulus diameter evaluated as percentage of the baseline diameter.¹⁸

Serum levels of soluble Nox2-derived peptide, a marker of NADPH oxidase activation, were detected by an ELISA method as previously described by Carnevale et al.¹⁹ Intra- and interassay coefficients of variation were 8.9% and 9%, respectively. Values are expressed as pg/mL.

Nitric oxide bioavailability was measured in serum by a colorimetric assay kit (Arbor Assays, Ann Arbor, MI). Intra- and interassay coefficients of variation were 4.4% and 6.8%, respectively. Values are expressed as $\mu\text{mol/L}$.

Quantification of isoprostanes was performed measuring serum 8-iso-PGF 2α -III (8-iso-PGF 2α -III) by enzyme immunoassay method (DRG International, Springfield, NJ). Intra- and interassay coefficients of variation were 5.8% and 5.0%, respectively. Values are expressed as pmol/L.

H $_2$ O $_2$ was evaluated by a colorimetric detection kit (Arbor Assays) and is expressed as $\mu\text{mol/L}$. Intra- and interassay coefficients of variation were 5.9% and 7.3%, respectively.

Serum H $_2$ O $_2$ breakdown activity (HBA) was measured with an HBA assay kit (Aurogene, Rome, Italy). Briefly, serum sample (2.5 μL) was incubated with H $_2$ O $_2$ and sample diluent at 37°C for 30 minutes. At the end of 30 minutes, stop solution was added and samples were then centrifuged at 3000 g for 5 minutes. Finally, the supernatant was read at 230 nm by a UV-visible single-beam spectrophotometer (A380; AOE Instruments Shanghai Co, Shanghai, China). Percentage of HBA was calculated according to the following formula: % Of HBA = $[(\text{Ac} - \text{As}) / \text{Ac}] \times 100$ where Ac is the absorbance of H $_2$ O $_2$ 1.4 mg/mL and As is the absorbance in the presence of the serum sample.

To measure vitamin E, serum samples (100 μL) were supplemented with tocopheryl acetate (internal standard), deproteinized by the addition of methanol, and centrifuged at 3000g for 10 minutes, as previously described.²⁰ The

collected upper phase was analyzed using an Agilent 1200 Infinity series high-performance liquid chromatography system (Agilent Technologies, Santa Clara, CA) equipped with an Eclipse Plus C18 column (4.6×100 mm). Vitamin E levels are presented as the ratio ($\mu\text{mol}/\text{mmol}$) between α -tocopherol concentration ($\mu\text{mol}/\text{L}$) and serum total cholesterol concentration (mmol/L).²¹

Platelet levels of soluble CD40L and P-selectin were measured in citrated blood samples centrifuged 15 minutes at 3000g. sCD40L levels were evaluated by a commercial immunoassay (Diacclone, Besancon, France), and values are expressed as ng/mL; intra- and interassay coefficients of variation were 4% and 6.8%, respectively. Soluble P-selectin levels were evaluated by a commercial immunoassay (Diacclone), and values are expressed as ng/mL; intra- and interassay coefficients of variation were 5.6% and 7.5%, respectively.

Serum levels of cotinine, a metabolite of nicotine and the primary biomarker for the determination of tobacco exposure,²² were evaluated by a commercial immunoassay (Origene, Rockville, MD), and values are expressed as ng/mL; intra- and interassay coefficients of variation were <10%.

Finally, we also measured systolic, diastolic, and mean blood pressure with an automated sphygmomanometer, averaging 3 measurements taken 3 minutes apart.

Notably, blood draws and blood pressure measurements were performed on the dominant arm (with blood draws preceding blood pressure measurements), whereas FMD was measured on the nondominant arm of each participant (Figure S1).

End Points and Analyses

End points were blood levels of soluble Nox2-derived peptide, nitric oxide (NO) bioavailability, H_2O_2 , HBA, 8-iso-PGF2 α -III, vitamin E, CD40 ligand, P-selectin, cotinine, FMD, blood pressure, and subjective satisfaction scores.

Statistical Analysis

Continuous variables are reported as means (SD) and categorical variables as counts (percentage). The main inferential analysis was based on a multilevel mixed-effects linear model, with an identity covariance matrix, forcing in the model each end point value, timing of sampling, product type, and their interaction as fixed effects, with participant and order as random effects (Data S1), similar to the model already used for the SUR-VAPES (Sapienza University of Rome-Vascular Assessment of Proatherosclerotic Effects of Smoking) 1 trial.¹² Additional analyses were run solely for descriptive purposes subgrouping by product type. Multilevel mixed-effects linear analyses are reported as point estimates

of effect (95% CIs) and corresponding *P* values. In addition, we performed nonparametric analyses with Wilcoxon signed-ranks test for satisfaction scores, with reporting based on median (first–third quartile). Although no formal sample-size computation was performed, the study was originally designed to include 20 nonsmokers and 20 smokers to achieve adequate statistical power, in keeping with a previous study from our group.¹² Because of issues in enrolling nonsmokers and have them use 3 different types of tobacco products, the nonsmoker arm of the study was discontinued at the beginning of the trial. Statistical significance was set at the 2-tailed 0.05 level, without multiplicity adjustment. Computations were performed with Stata software (version 13; StataCorp LP, College Station, TX).

Results

The study was conducted between September and December 2017. Notably, as many as 30 subjects had to be excluded after initial screening either because unwilling to refrain from smoking for the required washout period or for the concomitant presence of a clinical condition which qualified them for study exclusion. Table 1 shows baseline characteristics of study subjects: In particular, 70% were women and most usually smoked between 11 and 30 TCs per day. Moreover, 2 women were on oral contraceptives during the experiment.

Oxidative Stress Profile

To evaluate oxidative stress profile, we analyzed the levels of sNox2-dp, a small peptide released after platelet activation, which is a measure of Nox2 activation. As shown in Figure 1 and Tables 2 and 3, we observed an increase of sNox2-dp release after smoking each device (19.9 ± 9.9 versus 36.5 ± 6.8 pg/mL, $P<0.001$ for EVC; 23.1 ± 8.4 versus 44.1 ± 17.1 pg/mL, $P<0.001$ for TC; 22.8 ± 7.6 versus 29.9 ± 5.0 pg/mL, $P<0.001$ for HNBC; Table S1). Moreover, we evaluated the production of H_2O_2 , a nonradical oxygen species which permeates through cell membranes which is chemically stable. The results showed a significant increase of H_2O_2 for each smoking device (7.4 ± 3.4 versus 14.8 ± 2.9 $\mu\text{mol}/\text{L}$, $P<0.001$ for EVC; 7.6 ± 4.5 versus 19.5 ± 13.9 $\mu\text{mol}/\text{L}$, $P<0.001$ for TC; 6.3 ± 3.5 versus 12.8 ± 3.6 $\mu\text{mol}/\text{L}$, $P<0.001$ for HNBC; Figure S2; Tables 2 and 3).

Finally, we analyzed 8-iso-PGF2 α , an isoprostane produced by the nonenzymatic peroxidation of arachidonic acid in membrane phospholipids, which is a reliable biomarker of oxidative damage in vivo. We observed a significant increase of 8-iso-PGF2 α production after smoking (151 ± 18 versus 231 ± 31 , $P<0.001$ for EVC; 152 ± 20 versus 276 ± 29 , $P<0.001$ for TC; 158 ± 23 versus 207 ± 36 , $P<0.001$ for HNBC; Figure S3; Tables 2 and 3). Furthermore, as reported in

Table 1. Baseline Features

Subjects	20
Age, y	35±13
Male sex	6 (30%)
Height, cm	171±8
Weight, kg	70±18
Body mass index, kg/m ²	24±5
Waist-hip ratio	0.92±0.10
Time since beginning smoking, y	15±12
Cigarettes smoked per day	
<10	9 (45%)
11 to 20	9 (45%)
21 to 30	2 (10%)
Systolic blood pressure, mm Hg	116±15
Diastolic blood pressure, mm Hg	75±10
Total serum cholesterol, mg/dL	181±12
Family history of cardiovascular disease	10 (50%)
Previous cardiovascular disease	0
Concomitant pharmacological therapy (contraceptives, antiepileptics, or thyroid replacement therapy)	5 (25%)

Table 2, we observed that levels of oxidative stress biomarkers returned to baseline before each exposure given the adequate washout period.

Antioxidant Status

Antioxidant systems, which include enzymatic and nonenzymatic antioxidants, play a role in blocking harmful effects of reactive oxygen species. In this study, we evaluated levels of vitamin E, a powerful endogenous nonenzymatic antioxidant, and HBA, which is a measure of H₂O₂ neutralized by cellular enzymes. The results showed a significantly decrease of vitamin E levels after using EVC and TC (4.27±1.30 versus 2.71±1.07 μmol/mmol, *P*<0.001 for EVC; 3.95±1.62 versus 2.55±0.91 μmol/mmol, *P*<0.001 for TC; Figure S4; Tables 2 and 3). No changes were observed for vitamin E levels after using HNBC (4.11±1.09 versus 3.81±1.37 μmol/mmol, *P*=0.422; Figure S3; Tables 2 and 3). As regards HBA measurement, we observed a significant reduction in the ability to detoxify H₂O₂ after EVC and TC (54.5±18.4% versus 37.3±7.6%, *P*<0.001 for EVC; 54.1±17.1% versus 25.3±13.0%, *P*<0.001 for TC; Figure S5; Tables 2 and 3). A similarly significant reduction was observed for HBA after HNBC (55.4±9.9% versus 47.0±10.2%, *P*=0.006; Figure S3; Tables 2 and 3).

Biomarkers of antioxidant status returned to baseline levels before repeat exposure to each smoke device (Table 2).

Platelet Activation Assessment

To evaluate the role of different devices on platelet function, we analyzed 2 markers of platelet activation: sCD40L and soluble P-selectin. Results showed a significant increase in sCD40L levels with each device (3.20±1.16 versus 4.25±2.12 ng/mL, *P*=0.047 for EVC; 3.10±1.22 versus 5.26±1.97 ng/mL, *P*<0.001 for TC; 3.00±1.22 versus 4.18±1.56 ng/mL, *P*=0.007 for HNBC; Figure S6; Tables 2 and 3) and similar findings for soluble P-selectin (6.45±1.07 versus 7.97±1.65 ng/mL, *P*<0.001 for EVC; 6.76±1.28 versus 11.58±3.56 ng/mL, *P*<0.001 for TC; 6.63±1.22 versus 8.03±1.40 ng/mL, *P*<0.001 for HNBC; Figure S7; Tables 2 and 3). Biomarkers of platelet activation returned to baseline levels before repeat exposure to each smoke device (Table 2).

Endothelial Dysfunction

We evaluated the impact of smoking on endothelial dysfunction analyzing FMD, NO bioavailability, and blood pressure. FMD analysis showed a significantly reduction after using each device (6.14±3.17% versus 3.72±3.14%, *P*<0.001 for EVC; 6.20±3.26% versus 2.40±1.89%, *P*<0.001 for TC; 6.10±3.01% versus 3.79±2.68%, *P*<0.001 for HNBC; Figure 2; Tables 2 and 3). Similar results were observed for systolic blood pressure (121.7±6.5 versus 130.6±6.5 mm Hg, *P*<0.001 for EVC; 121.5±8.3 versus 132.4±6.2 mm Hg, *P*<0.001 for TC; 122.8±6.2 versus 129.7±6.4 mm Hg, *P*<0.001 for HNBC; Figure S8; Tables 2 and 3), diastolic blood pressure (72.2±4.4 versus 78.0±4.8 mm Hg, *P*<0.01 for EVC; 73.3±4.8 versus 80.2±5.2 mm Hg, *P*<0.01 for TC; 73.3±4.7 versus 76.9±5.0 mm Hg, *P*<0.01 for HNBC; Figure S9; Tables 2 and 3), and mean blood pressure (88.7±3.6 versus 95.5±3.6 mm Hg, *P*<0.001 for EVC; 89.4±4.7 versus 97.6±3.4 mm Hg, *P*<0.001 for TC; 89.8±4.4 versus 94.5±4.1 mm Hg, *P*<0.001 for HNBC; Figure S10; Tables 2 and 3).

Focusing on NO bioavailability, we observed a significant reduction after EVC and TC (24.9±13.6 versus 17.0±5.4 μmol/L, *P*=0.006 for EVC; 25.8±18.9 versus 12.7±6.6 μmol/L, *P*=0.006 for TC; Figure S11; Tables 2 and 3). Conversely, NO bioavailability did not change nonsignificantly with HNBCs (24.4±16.5 versus 19.8±6.6 μmol/L, *P*=0.206; Tables 2 and 3). Finally, parameters of endothelial dysfunction returned to baseline levels before each repeat exposure (Table 2).

Compliance Analysis

To evaluate smoking compliance, we analyzed serum levels of cotinine, a metabolite of nicotine and the primary biomarker for the determination of tobacco exposure, before and after




Figure 1. Impact of using electronic cigarette (EVC), traditional tobacco cigarette (TC), and heat-not-burn cigarette (HNBC) on blood levels of soluble Nox2-derived peptide. Box plots represent median, first quartile, third quartile, fifth percentile, 95th percentile, and outliers.

smoking. The results showed that all tobacco products increased significantly cotinine levels ($P<0.001$ for all; Tables 2 and 3).

Head-to-Head Comparisons of HNBC, EVC, and TC

HNBC and EVC were equally less impactful than TC for HBA, soluble CD40 ligand, and soluble P-selectin (both $P<0.05$ versus TC, $P>0.05$ for HNBC versus EVC). Similar trends were found for FMD ($P=0.872$ for HNBC versus EVC; $P=0.048$ for HNBC versus TC; $P=0.074$ for EVC versus TC), H_2O_2 ($P=0.522$ for HNBC versus EVC; $P=0.042$ for HNBC versus TC; $P=0.092$ for EVC versus TC), systolic blood pressure ($P=0.122$ for HNBC versus EVC; $P=0.002$ for HNBC versus TC; $P=0.121$ for EVC versus TC), diastolic blood pressure ($P=0.106$ for HNBC versus EVC; $P=0.046$ for HNBC versus TC; $P=0.532$ for EVC versus TC), and mean blood pressure ($P=0.053$ for HNBC versus EVC; $P=0.009$ for HNBC versus TC; $P=0.306$ for EVC versus TC). Conversely, HNBC were significantly less impactful than EVC and TC on soluble Nox2-derived peptide (respectively $P<0.001$ and $P=0.004$) and vitamin E (respectively $P=0.044$ and 0.018). Conversely, we found no nominally

significant interaction effects for NO bioavailability or cotinine.

Satisfaction Scores

Exploratory analysis of satisfaction scores (Table 4) suggested that TC were more enjoyable than both HNBC and EVC (both $P<0.05$; Figure S12). In addition, EVC proved less satisfying than both TC ($P=0.017$) and HNBC ($P=0.038$), without significant differences between these 2 ($P=0.239$; Figure S13). Similarly, EVC were rated as less likely to decrease desire for another cigarette than TC ($P=0.019$) and HNBC ($P=0.031$), without significant differences between these 2 ($P=0.581$; Figure S14).

Discussion

In the present cross-over randomized trial comparing the acute effects of HNBC, EVC, and TC, we found that use of any of these products was associated with acute detrimental effects on oxidative stress, antioxidant reserve, platelet function, FMD, and blood pressure.

Table 2. Descriptive Analysis*

Feature	Before Smoking			After Smoking		
	EVC	TC	HNBC	EVC	TC	HNBC
Soluble Nox2-derived peptide, pg/mL	19.9±9.9	23.1±8.4	22.8±7.6	36.5±6.8	44.1±17.1	29.9±5.0
Nitric oxide bioavailability, μmol/L	24.9±13.6	25.8±18.9	24.4±16.5	17.0±5.4	12.7±6.6	19.8±6.6
H ₂ O ₂ production, μmol/L	7.4±3.4	7.6±4.5	6.3±3.5	14.8±2.9	19.5±13.9	12.8±3.6
H ₂ O ₂ breakdown activity, %	54.5±18.4	54.1±17.1	55.4±9.9	37.3±7.6	25.3±13.0	47.0±10.2
8-iso-prostaglandin F-2α-III, pmol/L	151±18	152±20	158±23	231±31	276±29	207±36
Vitamin E, μmol/mmol	4.27±1.30	3.95±1.62	4.11±1.09	2.71±1.07	2.55±0.91	3.81±1.37
Soluble CD40 ligand, ng/mL	3.20±1.16	3.10±1.22	3.00±1.22	4.25±2.12	5.26±1.97	4.18±1.56
Soluble P-selectin, ng/mL	6.45±1.07	6.76±1.28	6.63±1.22	7.97±1.65	11.58±3.56	8.03±1.40
Flow-mediated dilation, %	6.14±3.17	6.20±3.26	6.10±3.01	3.72±3.14	2.40±1.89	3.79±2.68
Systolic blood pressure, mm Hg	121.7±6.5	121.5±8.3	122.8±6.2	130.6±6.5	132.4±6.2	129.7±6.4
Diastolic blood pressure, mm Hg	72.2±4.4	73.3±4.8	73.3±4.7	78.0±4.8	80.2±5.2	76.9±5.0
Mean blood pressure, mm Hg	88.7±3.6	89.4±4.7	89.8±4.4	95.5±3.6	97.6±3.4	94.5±4.1
Cotinine, ng/mL	31.6±16.6	34.4±19.3	30.4±12.0	64.4±11.1	65.5±10.2	61.0±16.7

EVC indicates electronic vaping cigarette; HNBC, heat-not-burn cigarette; TC, traditional tobacco cigarette.
 *Reported as mean±SD.

The validity of the chosen end points is well established. Indeed, systemic oxidative stress plays a fundamental role in vascular damage and in atherogenesis.²³ Reactive oxygen species are mostly generated by Nox2, an isoform from NADPH oxidase, that acts as an important regulator of

platelet-activation-associated thrombosis.²⁴ Indeed, Nox2 has been shown to be associated with several cardiovascular risk factors, such as hypercholesterolemia²⁵ and metabolic diseases,^{26,27} and in patients with peripheral artery disease.²⁸ Moreover, 8-iso-PGF2α and H₂O₂ production and endothelial

Table 3. Main Inferential Analysis*

Feature	Before vs After Smoking P Value			Interaction P Value		
	EVC	TC	HNBC	EVC vs TC	EVC vs HNBC	TC vs HNBC
Soluble Nox2-derived peptide	<0.001	<0.001	<0.001	0.318	0.004	0.001
Nitric oxide bioavailability	0.006	0.006	0.206	0.308	0.470	0.093
H ₂ O ₂ production	<0.001	<0.001	<0.001	0.092	0.522	0.042
H ₂ O ₂ breakdown activity	<0.001	<0.001	0.006	0.038	0.091	<0.001
8-iso-prostaglandin F-2α-III	<0.001	<0.001	<0.001	<0.001	0.004	<0.001
Vitamin E	<0.001	0.001	0.422	0.768	0.018	0.044
Soluble CD40 ligand	0.047	<0.001	0.007	0.046	0.849	0.071
Soluble P-selectin	<0.001	<0.001	<0.001	<0.001	0.821	<0.001
Flow-mediated dilation	<0.001	<0.001	<0.001	0.074	0.872	0.048
Systolic blood pressure	<0.001	<0.001	<0.001	0.121	0.122	0.002
Diastolic blood pressure	<0.001	<0.001	0.009	0.532	0.106	0.046
Mean blood pressure	<0.001	<0.001	<0.001	0.306	0.053	0.009
Cotinine	<0.001	<0.001	<0.001	0.782	0.722	0.935

EVC indicates electronic vaping cigarette; HNBC, heat-not-burn cigarette; TC, traditional tobacco cigarette.
 *Reported as P values, whereas point estimates of effect (95% CIs) are reported in Table S1.




Figure 2. Impact of using electronic cigarette (EVC), traditional tobacco cigarette (TC), and heat-not-burn cigarette (HNBC) on flow-mediated dilation (FMD). Box plots represent median, first quartile, third quartile, fifth percentile, 95th percentile, and outliers.

dysfunction are associated with cardiovascular events.^{28–31} Regarding physiological antioxidant capacity, the downregulation of the antioxidant system, such as vitamin E or enzymes devoted to H₂O₂ detoxification, is associated with an increased cardiovascular risk.^{32,33}

Focusing in detail on our results, a hierarchy of effects was apparent for some measures, with HNBC and EVC less impactful than TC on some dimensions of oxidative stress, antioxidant reserve, platelet function, and blood pressure. In addition, HNBC had less acute effects on soluble Nox2-derived peptide, 8-iso-PGF₂α-III, and vitamin E, and appeared more satisfying and capable of decreasing desire for continuing smoking than EVC.

The adverse impact of TC on health is well established, and smoking remains a leading preventable cause of morbidity and mortality worldwide.³⁴ Despite the lack of combustion and while less detrimental than TC, EVC cannot be considered harmless.^{9,12,35–38} The manufacturer of HNBC claims that they largely avoid pyrolysis and thus the generation of several harmful molecules, at least in relative terms, although this aspect may warrant additional study by independent

investigation.¹⁰ It is also very likely that the acute detrimental effects of HNBC are mediated, at least in part, by nicotine.^{39,40}

Our findings expand and refine knowledge on the acute comparative effects of HNBC, EVC, and TC and builds upon our recent work comparing EVC and TC for oxidative stress and FMD.¹² The fact that oxidative stress, platelet activation, and blood pressure are less impacted by HNBC and EVC than by TC suggests that they might be less detrimental, whereas some similarities between HNBC and TC, in terms of subjective satisfaction, might make HNBC more suitable than EVC as a risk-reduction product. Thus, these products cannot be considered equivalent in terms of risk, yet many opinion leaders, advocacy groups, and agencies call for complete abstinence from all products.⁴¹ Others have suggested that a more-pragmatic risk-reduction strategy could be more fruitful, at least in the short term.⁴² Our study provides information to guide future research on this topic, especially in light of the similar capacity of these products to deliver nicotine, as demonstrated by the nonsignificant interaction *P* values for cotinine.⁴³

Table 4. Satisfaction Scores*

Feature	Median (First–Third Quartile)			P Value		
	EVC	TC	HNBC	EVC vs TC	EVC vs HNBC	TC vs HNBC
Was the cigarette satisfying?	0 (0–50)	63 (38–75)	50 (25–75)	0.017	0.038	0.239
Was the cigarette enjoyable?	25 (0–50)	75 (50–88)	50 (25–75)	0.007	0.073	0.038
Did you experience a bad taste?	0 (0–50)	0 (0–50)	25 (0–75)	0.895	0.353	0.435
Did you suffer from unexpected cough?	0 (0–0)	0 (0–0)	0 (0–0)	0.970	0.361	0.305
Did smoking make you feel sick?	0 (0–0)	0 (0–0)	0 (0–0)	1	0.564	0.317
Did smoking give you energy?	0 (0–25)	0 (0–25)	0 (0–25)	0.882	0.910	0.471
Soon after smoking did your desire for another cigarette decrease?	25 (0–50)	75 (25–100)	75 (25–75)	0.019	0.031	0.581

EVC indicates electronic vaping cigarette; HNBC, heat-not-burn cigarette; TC, traditional tobacco cigarette.

*Scored on a subjective scale from 0 (no effect) to 100 (maximum effect).

Our work has, however, several limitations. The first is generalizability: we recruited from a homogeneous nonsmoking white population without overt clinical conditions, thus our findings cannot be directly applied to subjects with different features.⁴⁴ We eventually avoided the enrollment of nonsmokers in the study given difficulties in having them smoke 3 types of tobacco products, and also in order to avoid any possible incentive versus EVC or HNBC from the study findings. Indeed, the actual aim of the SUR-VAPES 2 trial is to provide preliminary data for subsequent longitudinal trials on the use of HNBC and EVC for smoking cessation in smokers with established cardiovascular disease. Conversely, we did not want to provide background for use of EVC or HNBC as a lower-risk surrogate of TC in nonsmokers. Indeed, exposure to EVC advertisements has been shown to cause increases in smoking impulses among adults who were former smokers or did not smoke at all, reduce teenagers' perceived risks of TC, and be associated with increased odds of EVC and TC use in both cross-sectional and longitudinal studies.^{45–47} Second, the small sample size and exposure period limited to 1 product per session prevent us from providing very precise effect estimates or understanding how strongly the observed trends in oxidative markers are correlated with smoking as opposed to statistical noise. Third, our focus on acute effects of a single smoking session of HNBC, EVC, and TC after an adequate washout cannot inform the chronic comparative impact of these products, and it remains unclear whether these chronic tobacco users may lead to more persistent vascular changes and biochemical profiles that do not represent the direct influence of each exposure. Accordingly, the study lacks longitudinal exposure and follow-up, decreasing our ability to understand whether the observed advantages of HNBC or EVC over TC from an oxidative standpoint are maintained with repetitive use of these smoking devices, or whether, for example, they lose significance in the long run. Fourth, we did not focus on other important health dimensions, in particular

those of pulmonary function and carcinogenesis, which require further investigation. Fifth, given the levels of cotinine at baseline assessments, compliance to washout recommendations was suboptimal. Nevertheless, this did not appear to affect between-device comparisons. Sixth, we did not adjust for multiplicity, but the mixed-effect design can be considered relatively robust for multiple inferential estimates. Nonetheless, the risk of type 1 error persists and should be borne in mind by readers. In addition, the mechanistic impact of HNBC and EVC on oxidative stress, FMD, and platelet function was not addressed in detail in our work, which was designed with a pragmatic scope.^{45,46} Details on satisfaction scores, while interesting given their important implications on the use of these approaches as a risk-reduction strategy, are limited by their exploratory scope and require additional analyses.^{47–49} Finally, the absence of a control group not receiving any tobacco product or an experimental group using a pure nicotine inhaler did not allow us to provide external baseline and repeated measures, nor focus specifically on the effect of selective nicotine administration on oxidative stress, FMD, and platelet activation.

In conclusion, while our findings should be regarded as preliminary and needing confirmation in other similar experiments, we found that use by a group of healthy adult smokers of HNBC, EVC, and TC, based on equivalent nicotine consumption, was associated with acute multidimensional adverse effects on a range of biological and physiological markers. The finding that HNBC is less impactful than both EVC and TC for some end points, while intriguing, requires further corroboration in larger studies with cohort design, long-term follow-up, and clinically relevant end points.

Author Contributions

Biondi-Zoccai, Sciarretta, Pignatelli, Carnevale, and Frati conceived and designed the study; Bullen participated in

study design, data interpretation, and manuscript drafting; Nocella and Cammisotto performed platelet analysis; Loffredo, Perri, and Pignatelli performed FMD assessment; De Falco, Carrizzo, and Chimenti performed NOX analysis; Marullo, Cavarretta, and Peruzzi performed patient enrollment; Valenti and Coluzzi provided questionnaires; Prati and Violi participated in data interpretation and manuscript drafting; Biondi-Zoccai performed statistical analysis; Biondi-Zoccai, Carnevale, Sciarretta, and Frati drafted the manuscript; all authors critically revised and approved the manuscript; Biondi-Zoccai acts as guarantor of the study.

Sources of Funding

This work was supported in part by grants from Sapienza University of Rome to Prof Frati (RM11715C7852D47E) and Prof De Falco (RM11715C7CCE6251), without any direct or indirect support from tobacco company.

Disclosures

Prof Biondi-Zoccai has consulted for Abbott Vascular and Bayer. The remaining authors have no disclosures to report.

References

- Walser T, Cui X, Yanagawa J, Lee JM, Heinrich E, Lee G, Sharma S, Dubinett SM. Smoking and lung cancer: the role of inflammation. *Proc Am Thorac Soc*. 2008;5:811–815.
- Stallones RA. The association between tobacco smoking and coronary heart disease. *Int J Epidemiol*. 2015;44:735–743.
- Carnevale R, Loffredo L, Pignatelli P, Nocella C, Bartimoccia S, Di Santo S, Martino F, Catasca E, Perri L, Violi F. Dark chocolate inhibits platelet isoprostanes via NOX2 down-regulation in smokers. *J Thromb Haemost*. 2012;10:125–132.
- Loffredo L, Carnevale R, Perri L, Catasca E, Augelletti T, Cangemi R, Albanese F, Piccheri C, Nocella C, Pignatelli P, Violi F. NOX2-mediated arterial dysfunction in smokers: acute effect of dark chocolate. *Heart*. 2011;97:1776–1781.
- Hahn J, Monakhova YB, Hengen J, Kohl-Himmelseher M, Schüssler J, Hahn H, Kuballa T, Lachenmeier DW. Electronic cigarettes: overview of chemical composition and exposure estimation. *Tob Induc Dis*. 2014;12:23.
- Tabuchi T, Kiyohara K, Hoshino T, Bekki K, Inaba Y, Kunugita N. Awareness and use of electronic cigarettes and heat-not-burn tobacco products in Japan. *Addiction*. 2016;111:706–713.
- Chen J, Bullen C, Dirks K. A comparative health risk assessment of electronic cigarettes and conventional cigarettes. *Int J Environ Res Public Health*. 2017;14:E382.
- Dinakar C, O'Connor GT. The health effects of electronic cigarettes. *N Engl J Med*. 2016;375:2608–2609.
- National Academies of Sciences, Engineering, and Medicine. *Public Health Consequences of E-Cigarettes*. Washington, DC: The National Academies Press; 2018. DOI: 10.17226/24952. <https://www.nap.edu/read/24952/chapter/1>. Accessed December 13, 2018.
- Schaller JP, Keller D, Poget L, Pratte P, Kaelin E, McHugh D, Cudazzo G, Smart D, Tricker AR, Gautier L, Yerly M, Reis Pires R, Le Bouhellec S, Ghosh D, Hofer I, Garcia E, Vanscheuwijck P, Maeder S. Evaluation of the Tobacco Heating System 2.2, part 2: chemical composition, genotoxicity, cytotoxicity, and physical properties of the aerosol. *Regul Toxicol Pharmacol*. 2016;81(Suppl 2):S27–S47.
- Auer R, Concha-Lozano N, Jacot-Sadowski I, Cornuz J, Berthet A. Heat-not-burn tobacco cigarettes: smoke by any other name. *JAMA Intern Med*. 2017;177:1050–1052.
- Carnevale R, Sciarretta S, Violi F, Nocella C, Loffredo L, Perri L, Peruzzi M, Marullo AG, De Falco E, Chimenti I, Valenti V, Biondi-Zoccai G, Frati G. Acute impact of tobacco vs electronic cigarette smoking on oxidative stress and vascular function. *Chest*. 2016;150:606–612.
- Nocella C, Biondi-Zoccai G, Sciarretta S, Peruzzi M, Pagano F, Loffredo L, Pignatelli P, Bullen C, Frati G, Carnevale R. Impact of tobacco versus electronic cigarette smoking on platelet function. *Am J Cardiol*. 2018;122:1477–1481.
- Hom S, Chen L, Wang T, Ghebrehiwet B, Yin W, Rubenstein DA. Platelet activation, adhesion, inflammation, and aggregation potential are altered in the presence of electronic cigarette extracts of variable nicotine concentrations. *Platelets*. 2016;27:694–702.
- Nabavizadeh P, Liu J, Havel CM, Ibrahim S, Derakhshandeh R, Jacob Iii P, Springer ML. Vascular endothelial function is impaired by aerosol from a single IQOS HeatStick to the same extent as by cigarette smoke. *Tob Control*. 2018;27:s13–s19.
- Shiffman S, Terhorst L. Intermittent and daily smokers' subjective responses to smoking. *Psychopharmacology*. 2017;234:2911–2917.
- Corretti MC, Anderson TJ, Benjamin EJ, Celermajer D, Charbonneau F, Creager MA, Deanfield J, Drexler H, Gerhard-Herman M, Herrington D, Vallance P, Vita J, Vogel R; International Brachial Artery Reactivity Task Force. Guidelines for the ultrasound assessment of endothelial-dependent flow-mediated vasodilation of the brachial artery: a report of the International Brachial Artery Reactivity Task Force. *J Am Coll Cardiol*. 2002;39:257–265.
- Loffredo L, Maccoccia A, Pignatelli P, Andreozzi P, Borgia MC, Cangemi R, Chiarrotti F, Violi F. Oxidative-stress-mediated arterial dysfunction in patients with peripheral arterial disease. *Eur Heart J*. 2007;28:608–612.
- Carnevale R, Silvestri R, Loffredo L, Novo M, Cammisotto V, Castellani V, Bartimoccia S, Nocella C, Violi F. Oleuropein, a component of extra virgin olive oil, lowers post-prandial glycaemia in healthy subjects. *Br J Clin Pharmacol*. 2018;84:1566–1574.
- Aust O, Sies H, Stahl W, Polidori MC. Analysis of lipophilic antioxidants in human serum and tissues: tocopherols and carotenoids. *J Chromatogr A*. 2001;936:83–93.
- Traber MG, Jialal I. Measurement of lipid-soluble vitamins—further adjustment needed? *Lancet*. 2000;355:2013–2014.
- Benowitz NL. Cotinine as a biomarker of environmental tobacco smoke exposure. *Epidemiol Rev*. 1996;18:188–204.
- Bernhard D, Wang XL. Smoking, oxidative stress and cardiovascular diseases—do anti-oxidative therapies fail? *Curr Med Chem*. 2007;14:1703–1712.
- Fuentes E, Gibbins JM, Holbrook LM, Palomo I. NADPH oxidase 2 (NOX2): a key target of oxidative stress-mediated platelet activation and thrombosis. *Trends Cardiovasc Med*. 2018;28:429–434.
- Loffredo L, Martino F, Carnevale R, Pignatelli P, Catasca E, Perri L, Calabrese CM, Palumbo MM, Baratta F, Del Ben M, Angelico F, Violi F. Obesity and hypercholesterolemia are associated with NOX2 generated oxidative stress and arterial dysfunction. *J Pediatr*. 2012;161:1004–1009.
- Del Ben M, Polimeni L, Carnevale R, Bartimoccia S, Nocella C, Baratta F, Loffredo L, Pignatelli P, Violi F, Angelico F. NOX2-generated oxidative stress is associated with severity of ultrasound liver steatosis in patients with non-alcoholic fatty liver disease. *BMC Gastroenterol*. 2014;14:81.
- Angelico F, Loffredo L, Pignatelli P, Augelletti T, Carnevale R, Pacella A, Albanese F, Mancini I, Di Santo S, Del Ben M, Violi F. Weight loss is associated with improved endothelial dysfunction via NOX2-generated oxidative stress down-regulation in patients with the metabolic syndrome. *Intern Emerg Med*. 2012;7:219–227.
- Loffredo L, Carnevale R, Cangemi R, Angelico F, Augelletti T, Di Santo S, Calabrese CM, Della Volpe L, Pignatelli P, Perri L, Basili S, Violi F. NOX2 up-regulation is associated with artery dysfunction in patients with peripheral artery disease. *Int J Cardiol*. 2013;165:184–192.
- Pignatelli P, Pastori D, Carnevale R, Farcomeni A, Cangemi R, Nocella C, Bartimoccia S, Vicario T, Saliola M, Lip GY, Violi F. Serum NOX2 and urinary isoprostanes predict vascular events in patients with atrial fibrillation. *Thromb Haemost*. 2015;113:617–624.
- Lambeth JD. NOX enzymes and the biology of reactive oxygen. *Nat Rev Immunol*. 2004;4:181–189.
- Violi F, Sanguigni V, Carnevale R, Plebani A, Rossi P, Finocchi A, Pignata C, De Mattia D, Martire B, Pietrogrande MC, Martino S, Gambineri E, Soresina AR, Pignatelli P, Martino F, Basili S, Loffredo L. Hereditary deficiency of gp91(phox) is associated with enhanced arterial dilatation: results of a multicenter study. *Circulation*. 2009;120:1616–1622.
- Carnevale R, Nocella C, Pignatelli P, Bartimoccia S, Stefanini L, Basili S, Novo M, D'Amico A, Cammisotto V, Pastori D, Violi F. Blood hydrogen peroxide break-down activity in healthy subjects and in patients at risk of cardiovascular events. *Atherosclerosis*. 2018;274:29–34.

33. Vardi M, Levy NS, Levy AP. Vitamin E in the prevention of cardiovascular disease: the importance of proper patient selection. *J Lipid Res.* 2013;54:2307–2314.
34. Jha P, Peto R. Global effects of smoking, of quitting, and of taxing tobacco. *N Engl J Med.* 2014;370:60–68.
35. Goniewicz ML, Knysak J, Gawron M, Kosmider L, Sobczak A, Kurek J, Prokopowicz A, Jablonska-Czapla M, Rosik-Dulewska C, Havel C, Jacob P III, Benowitz N. Levels of selected carcinogens and toxicants in vapour from electronic cigarettes. *Tob Control.* 2014;23:133–139.
36. Moheimani RS, Bhetraratana M, Peters KM, Yang BK, Yin F, Gornbein J, Araujo JA, Middlekauff HR. Sympathomimetic effects of acute E-cigarette use: role of nicotine and non-nicotine constituents. *J Am Heart Assoc.* 2017;6:e006579. DOI: 10.1161/JAHA.117.006579.
37. Moheimani RS, Bhetraratana M, Yin F, Peters KM, Gornbein J, Araujo JA, Middlekauff HR. Increased cardiac sympathetic activity and oxidative stress in habitual electronic cigarette users: implications for cardiovascular risk. *JAMA Cardiol.* 2017;2:278–284.
38. Glasser AM, Collins L, Pearson JL, Abudayyeh H, Niaura RS, Abrams DB, Villanti AC. Overview of electronic nicotine delivery systems: a systematic review. *Am J Prev Med.* 2017;52:e33–e66.
39. Liu X, Lugo A, Spizzichino L, Tabuchi T, Pacifici R, Gallus S. Heat-not-burn tobacco products: concerns from the Italian experience. *Tob Control.* 2019;28:113–114.
40. Davis B, Williams M, Talbot P. iQOS: evidence of pyrolysis and release of a toxicant from plastic. *Tob Control.* 2019;28:34–41.
41. Avdalovic MV, Murin S. POINT: does the risk of electronic cigarettes exceed potential benefits? Yes. *Chest.* 2015;148:580–582.
42. Middlekauff HR. COUNTERPOINT: does the risk of electronic cigarettes exceed potential benefits? No. *Chest.* 2015;148:582–584.
43. Fairchild AL, Lee JS, Bayer R, Curran J. E-cigarettes and the harm-reduction continuum. *N Engl J Med.* 2018;378:216–219.
44. Mastrangeli S, Carnevale R, Cavarretta E, Sciarretta S, Peruzzi M, Marullo AGM, De Falco E, Chimenti I, Valenti V, Bullen C, Roeber L, Frati G, Biondi-Zoccai G. Predictors of oxidative stress and vascular function in an experimental study of tobacco versus electronic cigarettes: a post-hoc analysis of the SUR-VAPES 1 (Sapienza University of Rome-Vascular Assessment of Proatherosclerotic Effects of Smoking) Study. *Tob Induc Dis.* 2018;16:11.
45. Calvieri C, Tanzilli G, Bartimoccia S, Cangemi R, Arrivi A, Dominici M, Cammisotto V, Viceconte N, Mangieri E, Frati G, Violi F. Interplay between oxidative stress and platelet activation in coronary thrombus of STEMI patients. *Antioxidants (Basel).* 2018;7:E83.
46. Carnevale R, Loffredo L, Nocella C, Bartimoccia S, Bucci T, De Falco E, Peruzzi M, Chimenti I, Biondi-Zoccai G, Pignatelli P, Violi F, Frati G. Epicatechin and catechin modulate endothelial activation induced by platelets of patients with peripheral artery disease. *Oxid Med Cell Longev.* 2014;2014:691015.
47. Biondi-Zoccai G, Peruzzi M, Frati G. E-cigarettes, incentives, and drugs for smoking cessation. *N Engl J Med.* 2018;379:991–992.
48. Auf R, Trepka MJ, Selim M, Ben Taleb Z, De La Rosa M, Cano MÁ. E-cigarette marketing exposure and combustible tobacco use among adolescents in the United States. *Addict Behav.* 2018;78:74–79.
49. King AC, Smith LJ, McNamara PJ, Cao D. Second generation electronic nicotine delivery system vape pen exposure generalizes as a smoking cue. *Nicotine Tob Res.* 2018;20:246–252.

SUPPLEMENTAL MATERIAL

Data S1.

Additional details on statistical analysis

Analysis was based on a long-formatted database, as the sample below:

ID	Order	Product	Timing	Measurement
1	1	A	Baseline	Y1
1	2	A	Post	Y2
1	3	B	Baseline	Y3
1	4	B	Post	Y4
1	5	C	Baseline	Y5
1	6	C	Post	Y6
2	1	B	Baseline	Y7
2	2	B	Post	Y8
2	3	C	Baseline	Y9
2	4	C	Post	Y10
2	5	A	Baseline	Y11
2	6	A	Post	Y12
...

The model for the within-product before-after analysis was:

$$Y_{ij} = \beta_0 + \beta_1 \text{Time} + u_1 \text{ID} + u_2 \text{Order} + \varepsilon$$

with β representing the fixed effects, u the random effects, and ε the Gaussian error term.

The Stata code for the within-product before-after analysis was:

```
mixed Measurement Timing || ID:, || Order:, covariance(identity)
```

Conversely, the model for the between-product interaction analysis was:

$$Y_{ij} = \beta_0 + \beta_1 \text{Time} + \beta_2 \text{Product} + \beta_3 \text{Time} * \text{Product} + u_1 \text{ID} + u_2 \text{Order} + \varepsilon$$

with β representing the fixed effects, u the random effects, and ε the Gaussian error term.

Accordingly, the Stata code for the between-product interaction analysis was:

```
mixed Measurement Timing##Product || ID:, || Order:, covariance(identity)
```


A sample screenshot for the Stata output is the following:

```
. mixed FMD Pre0Post1##TC1EC2HNS3 || ID:, || Order:, covariance(identity)
```

Performing EM optimization:

Performing gradient-based optimization:

```
Iteration 0: log likelihood = -248.25534
Iteration 1: log likelihood = -246.75537
Iteration 2: log likelihood = -246.71072
Iteration 3: log likelihood = -246.71072
```

Computing standard errors:

```
Mixed-effects ML regression           Number of obs   =       115
```

Group Variable	No. of Groups	Observations per Group		
		Minimum	Average	Maximum
ID	20	3	5.8	6
Order	58	1	2.0	2

```
Wald chi2(5) = 91.50
Prob > chi2 = 0.0000
Log likelihood = -246.71072
```

FMD	Coef.	Std. Err.	z	P> z	[95% Conf. Interval]	
1.Pre0Post1	-3.8114	.5404618	-7.05	0.000	-4.870685	-2.752114
TC1EC2HNS3						
2	-.0707635	.5545494	-0.13	0.898	-1.15766	1.016133
3	-.1178998	.5404618	-0.22	0.827	-1.177185	.9413859
Pre0Post1#TC1EC2HNS3						
1 2	1.389178	.7785131	1.78	0.074	-.1366801	2.915035
1 3	1.5004	.7580794	1.98	0.048	.0145915	2.986208
_cons	6.2149	.6280043	9.90	0.000	4.984034	7.445766

Random-effects Parameters	Estimate	Std. Err.	[95% Conf. Interval]	
ID: Identity				
var(_cons)	4.871662	1.691218	2.467056	9.620007
Order: Identity				
var(_cons)	1.25e-17	6.91e-17	2.56e-22	6.14e-13
var(Residual)	2.825853	.4095205	2.127131	3.754094

```
LR test vs. linear regression:      chi2(2) = 70.26   Prob > chi2 = 0.0000
```

Note: LR test is conservative and provided only for reference.

Table S1. Additional inferential analysis.

Feature	Point estimate of effect (95% confidence interval)		
	EVC vs TC	EVC vs HNBC	TC vs HNBC
Primary endpoints			
Soluble Nox2-derived peptide (pg/mL)	-4.30 (-12.74; 4.14)	-9.55 (-16.00; -3.10)	-13.85 (-22.29; -5.41)
Flow-mediated dilation (%)	1.39 (-0.14; 2.92)	0.11 (-1.25; 1.47)	1.50 (0.02; 2.97)
Additional endpoints			
Nitric oxide bioavailability (μM)	5.18 (-4.77; 15.13)	3.35 (-5.73; 12.42)	8.52 (-1.42; 18.48)
H ₂ O ₂ production (μmol/L)	-4.58 (-9.89; 0.74)	-0.94 (-3.81; 1.94)	-5.51 (-10.83; -0.20)
H ₂ O ₂ breakdown activity (%)	11.63 (0.64; 22.61)	8.76 (-1.40; 18.91)	20.38 (9.40; 31.37)
8-iso-prostaglandin F-2α-III (pmol/L)	-44.4 (-66.1; -22.7)	-31.0 (-52.2; -9.8)	-75.4 (-97.1; -53.7)
Vitamin E (μmol/mmol)	1.09 (0.03; 2.15)	1.25 (0.22; 2.29)	-0.16 (-1.22; 0.90)
Soluble CD40 ligand (ng/mL)	-1.38 (-2.74; -0.02)	0.13 (-1.21; 1.47)	-1.25 (-2.61; 0.11)
Soluble P-selectin (ng/ml)	-3.30 (-4.92; -1.68)	-0.13 (-1.21; 0.96)	-3.43 (-5.04; -1.81)
Systolic blood pressure (mm Hg)	-2.05 (-4.64; 0.54)	-1.95 (-4.42; 0.52)	-4.00 (-6.59; -1.41)
Diastolic blood pressure (mm Hg)	-1.00 (-4.14; 2.24)	-2.20 (-4.87; 0.47)	-3.20 (-6.34; -0.06)
Mean blood pressure (mm Hg)	-1.35 (-3.93; 1.23)	-2.12 (-4.26; 0.03)	-3.47 (-6.05; -0.88)
Cotinine (ng/mL)	1.70 (-10.32; 13.72)	-2.20 (-14.31; 9.91)	-0.50 (-12.52; 11.52)

EVC=electronic vaping cigarette; H₂O₂=hydrogen peroxide; HNBC=heat-not-burn cigarette; TC=traditional tobacco cigarette

Figure S1. Consolidated Standards of Reporting Trials (CONSORT) subject flow diagram (left panel), and measurement protocol (right panel). BP=blood pressure; EVC=electronic vaping cigarette; FMD=flow-mediated dilation; HNBC=heat-not-burn cigarette; TC=traditional tobacco cigarette.




Figure S2. Impact of using electronic vaping cigarette (EVC), traditional tobacco cigarette (TC), and heat-not-burn cigarette (HNBC) on hydrogen peroxide (H₂O₂) production. Boxplots represent median, 1st quartile, 3rd quartile, 5th percentile, 95th percentile, and outliers.




Figure S3. Impact of using electronic vaping cigarette (EVC), traditional tobacco cigarette (TC), and heat-not-burn cigarette (HNBC) on 8-iso-prostaglandin F-2 α -III (8-iso-PGF2a). Boxplots represent median, 1st quartile, 3rd quartile, 5th percentile, 95th percentile, and outliers.




Figure S4. Impact of using electronic vaping cigarette (EVC), traditional tobacco cigarette (TC), and heat-not-burn cigarette (HNBC) on vitamin E.

Boxplots represent median, 1st quartile, 3rd quartile, 5th percentile, 95th percentile, and outliers.




Figure S5. Impact of using electronic vaping cigarette (EVC), traditional tobacco cigarette (TC), and heat-not-burn cigarette (HNBC) on hydrogen peroxide (H₂O₂) breakdown activity (HBA). Boxplots represent median, 1st quartile, 3rd quartile, 5th percentile, 95th percentile, and outliers.




Figure S6. Impact of using electronic vaping cigarette (EVC), traditional tobacco cigarette (TC), and heat-not-burn cigarette (HNBC) on soluble CD40 ligand. Boxplots represent median, 1st quartile, 3rd quartile, 5th percentile, 95th percentile, and outliers.




Figure S7. Impact of using electronic vaping cigarette (EVC), traditional tobacco cigarette (TC), and heat-not-burn cigarette (HNBC) on soluble P-selectin. Boxplots represent median, 1st quartile, 3rd quartile, 5th percentile, 95th percentile, and outliers.




Figure S8. Impact of using electronic vaping cigarette (EVC), traditional tobacco cigarette (TC), and heat-not-burn cigarette (HNBC) on systolic blood pressure (SBP). Boxplots represent median, 1st quartile, 3rd quartile, 5th percentile, 95th percentile, and outliers.




Figure S9. Impact of using electronic vaping cigarette (EVC), traditional tobacco cigarette (TC), and heat-not-burn cigarette (HNBC) on diastolic blood pressure (DBP). Boxplots represent median, 1st quartile, 3rd quartile, 5th percentile, 95th percentile, and outliers.




Figure S10. Impact of using electronic vaping cigarette (EVC), traditional tobacco cigarette (TC), and heat-not-burn cigarette (HNBC) on mean blood pressure (MBP). Boxplots represent median, 1st quartile, 3rd quartile, 5th percentile, 95th percentile, and outliers.




Figure S11. Impact of using electronic vaping cigarette (EVC), traditional tobacco cigarette (TC), and heat-not-burn cigarette (HNBC) on nitric oxide (NO) bioavailability. Boxplots represent median, 1st quartile, 3rd quartile, 5th percentile, 95th percentile, and outliers.




Figure S12. Impact of using electronic vaping cigarette (EVC), traditional tobacco cigarette (TC), and heat-not-burn cigarette (HNBC) on smoking satisfaction, appraised with the explicit question “Was the cigarette enjoyable?”, and answers scored using a subjective scale from 0 (no effect) to 100 (maximum effect). Boxplots represent median, 1st quartile, 3rd quartile, 5th percentile, 95th percentile, and outliers.




Figure S13. Impact of using electronic vaping cigarette (EVC), traditional tobacco cigarette (TC), and heat-not-burn cigarette (HNBC) on smoking satisfaction, appraised with the explicit question “Was the cigarette satisfying?”, and answers scored using a subjective scale from 0 (no effect) to 100 (maximum effect). Boxplots represent median, 1st quartile, 3rd quartile, 5th percentile, 95th percentile, and outliers.




Figure S14. Impact of using electronic vaping cigarette (EVC), traditional tobacco cigarette (TC), and heat-not-burn cigarette (HNBC) on smoking satisfaction, appraised with the explicit question “Soon after smoking did your desire for another cigarette decrease?”, and answers scored using a subjective scale from 0 (no effect) to 100 (maximum effect). Boxplots represent median, 1st quartile, 3rd quartile, 5th percentile, 95th percentile, and outliers.

